REFERENCES:

1. Adriano R, Vitale C and Marangella M (2000). Epidemiology of nephrolithiasis. J Nephro 13: 65-70.
2. Agarwal R, Gupta SK, Agrawal SS, Srivastava S and Saxena R (2008). Oculohypotensive effects of *Foeniculum vulgare* in Experimental models of glaucoma. Indian J Physiol Pharmacol 52(1): 77-83.
3. Alexandrovich I, Rakovitskaya O, Kolmo E, Tatyana S and Shushunov S (2003). The effect of fennel (*Foeniculum vulgare*) seed oil emulsion in infantile colic: A randomized, placebo-controlled study. Alternative Therapies 9(4): 58-61.
4. Al-Kohlany KM, Shokeir AA, Mosbah A, Mohsen T, Shoma AM, Eraky I, El-Kenawy M and El-Kappany HA (2005). Treatment of complete staghorn stones: a prospective randomized comparison of open surgery versus percutaneous nephrolithotomy. J Urol. 173(2): 469-473.
5. Bailly GG, Norman RW and Thompson C (2000). Effects of dietary fat on the urinary risk factors of calcium stone disease. Urology 56: 40-44.
6. Barros ME, Lima R, Mercuri LP, Matos JR, Schor N and Boim MA (2006). Effect of extract of *Phyllanthus niruri* on crystal deposition in experimental urolithiasis. Urol Res 34: 351-357.
7. Basavaraj DR, Biyani CS, Browning AJ and Cartledge JJ (2007). The Role of Urinary Kidney Stone Inhibitors and Promoters in the Pathogenesis of Calcium Containing Renal Stones. EAU-EBU Update Series 5: 126-136.
8. Bhatt PA and Paul P (2008). Analysis of urinary stone constituents using powder X-ray diffraction and FT–IR. J. Chem. Sci. 120(2): 267–273.
9. Bihl G and Meyers A (2001). Recurrent renal stone disease-advances in pathogenesis and clinical management. Lancet 358: 651-656.
10. Bijarnia RK, Kaur T, Singla SK and Tandon C (2010). Non-surgical management therapies for kidney stones. J Pharm Educ Res.: 21-24.
11. Birdane FM, Cemek M, Birdane YO, Gulcin I and Buyukokuroglu ME (2007). Beneficial effects of *Foeniculum vulgare* on ethano-induced acute gastric mucosal injury in rats. World J Gastroentero 13(4): 607-611.
12. Choi EM, Hwang JK (2004). Antiinflammatory, analgesic and antioxidant activities of the fruit of *Foeniculum vulgare*. Fitoterapia 75: 557– 565.

13. Curhan GC, Willett WC, Speizer FE, Spiegelman D and Stampfer MJ (1997). Comparison of Dietary Calcium with Supplemental Calcium and Other Nutrients as Factors Affecting the Risk for Kidney Stones in Women. Annals of Internal Medicine 126(7): 497-504.
14. Doddola S, Pasupulati H, Koganti B and Prasad KVSRG (2008). Evaluation of *Sesbania grandiflora* for antiurolithiatic and antioxidant properties. J Nat Med 62: 300-307.
15. Duarte RJ, Mitre AI, Denes FT, Giron AM, Koch VH and Arap S (2002). Extracorporeal lithotripsy for the treatment of urolithiasis in children. Journal de Pediatria. 78(5): 367-370.
16. Ebeed NM, Abdou HS, Booles HF, Salah SH, Ahmed ES and Fahmy K (2010). Antimutagenic and Chemoprevention Potentialities of Sweet Fennel (*Foeniculum vulgare* Mill.) Hot Water Crude Extract. Journal of American Science 6(9): 831-842.
17. Galib BS, Behera and Kar AC (2006). Lithotriptic activity of swadamstradi kwatha seadamstradi chan Vati-an experimental evaluation. JDRAS 27: 8-18.
18. Goldfarb DS and Coe FL (1999). Beverages, Diet and Prevention of Kidney Stones. American Journal of Kidney Diseases 33(2): 398-403.
19. Hall PM (2009). Nephrolithiasis: Treatment, causes and prevention. Cleveland Clinic Journal of Medicine 79(10): 583-591.
20. Harrison L (2004). Principle of international medicine. Mc graw hill. 2(15): 1615-1620.
21. Hughes P (2007). Kidney stones epidemiology. Nephrology 12: S26–S30.
22. Iqbal S, Ahmed R and Muhammad F (1999). Evaluation of Diuretic Effect of an Indigenous Herbal Preparation in Goats. International Journal of Agriculture & Biology 1(3): 136-137.
23. Javidnia K, Dastgheib L, Samani SM and Nasiri A (2003). Antihirsutism activity of Fennel (fruits of *Foeniculum vulgare*) extract A double-blind placebo controlled study. Phytomedicine 10: 455-458.
24. Joseph KC, Parekh BB and Joshi MJ (2005). Inhibition of growth of urinary type calcium hydrogen phosphate dihydrate crystals by tartaric acid and tamarind. Current Science 88(8): 1232-1238.

25. Karadi RV, Gadge NB, Alagawadi KR and Savadi RV (2006). Effect of *Moringa oleifera* Lam. root-wood on ethylene glycol induced urolithiasis in rats. Journal of Ethnopharmacology 105: 306-311.
26. Kaur GJ and Arora DS (2009). Antibacterial and phytochemical screening of *Anethum graveolens*, *Foeniculum vulgare* and *Trachyspermum ammi.* BMC Complementary and Alternative Medicine 30(9): 1-10.
27. Knoll T (2007). Stone Disease. European Urology Supplements 6: 717-722.
28. Knoll T (2010). Epidemiology, Pathogenesis and Pathophysiology of Urolithiasis. European Urology Supplements 9: 802-806.
29. Lee YH, Tsai JY and Haung JK (2000). Combined use of 30% Lactose rich diet and 1 % ethylene glycol: A new animal model for study of urolithiasis. J Uro ROC 11: 149-154.
30. Oka Y, Nacar S, Putievsky E, Ravid U, Yaniv Z and Spiegel Y (2000). Nematicidal activity of essential oils and their components against the root knot nematode. Phytopathology 90(7): 710-715.
31. Ostad SN, Soodi M,Shariffzadeh M, Khorshidi N, Marzban H (2001). The effect of fennel essential oil on uterine contraction as a model for dysmenorrheal, pharmacology and toxicology study. Journal of Ethnopharmacology 76: 299-304.
32. Ozbek H (2005). The anti-inflammatory activity of the *Foeniculum vulgare* L. essential oil and investigation of its median lethal dose in rats and mice. International Journal of Pharmacology 1(4): 329-331.
33. Ozbek H, Bayram I, Ugras S and Cengiz N (2006). Investigation of hepatoprotective effect of *Foeniculum vulgare* fixed oil in rats. Research Journal of Medicine and Medical Sciences 1(2): 72-76.
34. Ozbek H, Ugras S, Bayram I, Uygan I, Erdogan E, Ozturk A and Huyut Z (2004). Hepatoprotective effect of *Foeniculum vulgare* essential oil: A carbon-tetrachloride induced liver fibrosis model in rats. Scand. J. Lab. Anim. Sci. 31(1): 9-17.
35. Pai MBH, Prashant GM, Murlikrishna KS, Shivakumar KM and Chandu GN (2010). Antifungal efficacy of *Punica granatum, Acacia nilotica, Cuminum cyminum* and *Foeniculum vulgare* on *Candida albicans*: An in vitro study. Indian J Dent Res 21(3): 334-336.

36. Pais VM, Lowe G, Lallas CD, Preminger GM and Assimos DG (2006). Xanthine Urolithiasis. Urology 67(5): 1084.e9-1084.e11.
37. Pak CYC (1998). Kidney Stones. Lancet 351: 1797-1801.
38. Pradhan M, Sribhuwaneswari S, Karthikeyan D, Minz S, Sure P, Chandu An, Mishra U, Kamalakannan K, Saravanankumar A and Sivakumar T (2008). In-vitro Cytoprotection Activity of *Foeniculum vulgare* and *helicteres isora* in Cultured Human Blood Lymphocytes and Antitumor Activity against B16F10 Melanoma Cell Line. Research J Pharm and Tech 1(4): 450-452.
39. Sarica K, Altay B and Erturhan S (2008). Effect of Being Overweight on Stone-Forming Risk Factors. Urology 71: 771-775.
40. Sas DJ, Hursey TC, Shatat IF and Orak JK (2010). Increasing Incidence of kidney Stones in Children Evaluated in the Emergency Department. J Pediatr 157: 132-137.
41. Sharma AP and Filler G (2010). Epidemiology of pediatric urolithiasis. Indian Journal of Urology 26(4): 516-522.
42. Shukla HS, Dubey P and Chaturvedi RV (1989). Antiviral properties of essential oils of *Foeniculum vulgare* and *Pimpinella anisum L.* Agronomie 9: 277-279.
43. Tiselius HG (2000). Stone Incidence and Prevention. Brazilian Journal of Urology 26(5): 452-462.
44. Tognolini M, Ballabeni V, Bertoni S, Bruni R, Impicciatore M, Barocelli E (2007). Protective effect of *Foeniculum vulgare* essential oil and anethole in an experimental model of thrombosis. Pharmacological Research 56: 254-260.
45. Venkataranganna MV, Anturlikar SD, Gopumadhavan S, Prakash NS, Rafiq M, Murthy GS and Mitra SK (2002). Antispasmodic activity of SJ-200 (Himcospaz) An herbal preparation. Pharmaceutical Biology 40(6): 416-421.
46. Wasserstaein AG (2005). Nephrolithiasis. American Journal of Kidney Diseases 45(2): 422-428.
47. Wasserstein A (2005). Core curriculum in nephrology. American journal of Kidney disease. 42(2): 422-428.
48. Yasar F, Kusvuran S and Ellialtioglu S (2006). Determination of anti-oxidant activities in some melon (*Cucumis melo* L.) varieties and cultivars under salt stress. Journal of Horticultural Science & Biotechnology 81(4): 627-630.